

**Class X  
(CBSE 2019)  
Mathematics  
Abroad (Set-2)**

---

Please keep a pen and paper ready for rough work

JavaScript must be enabled on your browsers

Your time countdown will begin as soon as you click the "Proceed/Resume" button

Your test will automatically end when the allotted time for the test runs out

If the browser window closes during the test, then you can resume the test from Test Papers page

During a test, you can move backwards and forwards and jump to any question you like

You can change the answers to your questions

You will see how many questions you have attempted, answered and skipped on the right side

---

**Question 1**

For what values of  $k$  does the quadratic equation  $4x^2 - 12x - k = 0$  have no real roots?

**SOLUTION:**

We have been given the quadratic equation:

$$4x^2 - 12x - k = 0$$

To have no real roots means discriminant should be less than zero.

$$D = b^2 - 4ac$$

$$b^2 - 4ac < 0$$

Plugging the values in the formula of discriminant

$$(-12)^2 - 4(4)(-k) < 0$$

$$144 + 16k < 0$$

$$k < -9$$

Therefore, for  $k < -9$  the quadratic equation will have no real roots.

Question 2

Find the distance between the points  $(a, b)$  and  $(-a, -b)$ .

**SOLUTION:**

Using distance formula:

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

Here,  $x_1 = a$ ,  $y_1 = b$ ,  $x_2 = -a$  and  $y_2 = -b$

On substituting the values in the formula we get

$$\sqrt{(-a - a)^2 + (-b - b)^2}$$

$$= \sqrt{(-2a)^2 + (-2b)^2}$$

$$= \sqrt{4a^2 + 4b^2}$$

$$= 2\sqrt{a^2 + b^2}$$

Therefore, the distance between  $(a, b)$  and  $(-a, -b)$  is  $2\sqrt{a^2 + b^2}$

Question 3

Find a rational number between  $\sqrt{2}$  and  $\sqrt{7}$ .

OR

Write the number of zeroes in the end of a number whose prime factorization is  $2^2 \times 5^3 \times 3^2 \times 17$ .

**SOLUTION:**

We know

$$\sqrt{2} = 1.414$$

$$\sqrt{7} = 1.732$$

So, rational number between  $\sqrt{2}$  and  $\sqrt{7}$  will be  $1.5 = \frac{3}{2}$ .

OR

Given prime factorisation is  $2^2 \times 5^3 \times 3^2 \times 17$ .

A number will have zero at the end when we have  $2 \times 5$ .

In  $2^2 \times 5^3 \times 3^2 \times 17$  we will have 2 zeroes as  $(2^2 \times 5^2) \times 5 \times 3^2 \times 17$ .

Question 4

Let  $\triangle ABC \sim \triangle DEF$  and their areas be respectively,  $64 \text{ cm}^2$  and  $121 \text{ cm}^2$ . If  $EF = 15.4 \text{ cm}$ , find  $BC$ .

SOLUTION:

Given:  $\triangle ABC \sim \triangle DEF$

We know ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

$$\frac{\text{ar}\triangle ABC}{\text{ar}\triangle DEF} = \left(\frac{BC}{EF}\right)^2$$

$$\Rightarrow \frac{64}{121} = \left(\frac{BC}{15.4}\right)^2$$

$$\Rightarrow \left(\frac{8}{11}\right)^2 = \left(\frac{BC}{15.4}\right)^2$$

$$\Rightarrow \frac{8}{11} = \frac{BC}{15.4}$$

$$\Rightarrow BC = \frac{8 \times 15.4}{11} = 11.2 \text{ cm}$$

Thus,  $BC = 11.2 \text{ cm}$ .

Question 5

Evaluate:

$$\frac{\tan 65^\circ}{\cot 25^\circ}$$

OR

Express  $(\sin 67^\circ + \cos 75^\circ)$  in terms of trigonometric ratios of the angle between  $0^\circ$  and  $45^\circ$ .

SOLUTION:

$$\frac{\tan 65^\circ}{\cot 25^\circ}$$

$$= \frac{\tan(90^\circ - 25^\circ)}{\cot 25^\circ} \quad (\because \tan(90^\circ - \theta) = \cot \theta)$$

$$= \frac{\cot 25^\circ}{\cot 25^\circ}$$

$$= 1$$

OR

$$(\sin 67^\circ + \cos 75^\circ)$$

$$= (\sin(90^\circ - 23^\circ) + \cos(90^\circ - 25^\circ)) \quad (\because \sin(90^\circ - \theta) = \cos \theta \text{ and } \cos(90^\circ - \theta) = \sin \theta)$$

$$= (\cos 23^\circ + \sin 25^\circ)$$

Question 6

Find the number of terms in the A.P. :  $18, 15\frac{1}{2}, 13, \dots, -47$ .

**SOLUTION:**

We have been given an A.P

$18, 15\frac{1}{2}, 13, \dots -47$

Here,  $a = 18, d = 15\frac{1}{2} - 18 = \frac{-5}{2}, a_n = -47$

We will find  $n$  using

$$a_n = a + (n - 1)d$$

Plugging the values in the formula we get:

$$-47 = 18 + (n - 1) \left( \frac{-5}{2} \right)$$

$$-47 = 18 - \frac{5}{2}n + \frac{5}{2}$$

$$n = 27$$

Therefore, there are 27 terms in an A.P

Question 7

A bag contains 15 balls, out of which some are white and the others are black. If the probability of drawing a black ball at random from the bag is  $\frac{2}{3}$ , then find how many white balls are there in the bag.

**SOLUTION:**

Total number of balls 15

Probability of drawing a black ball at random is  $\frac{2}{3}$

probability of black ball + probability of white ball = 1

Probability of white ball = 1 - probability of black ball

$$\text{Probability of drawing a white ball} = 1 - \frac{2}{3} = \frac{1}{3}$$

$$\text{Therefore, number of white balls} = 15 \times \frac{1}{3} = 5$$

Question 8

A card is drawn at random from a pack of 52 playing cards. Find the probability of drawing a card which is neither a spade nor a king.

**SOLUTION:**

We have total number of cards 52

And in deck of 52 cards number of spade are 13

And number of king = 4

But, out of these 4 kings, 1 king is already included in 13 spades card.

So, we will remove all the spade and king that is  $52 - (13+3) = 36$

Therefore, probability of neither a spade nor a king is  $\frac{36}{52} = \frac{9}{13}$

### Question 9

Find the solution of the pair of equation :

$$\frac{3}{x} + \frac{8}{y} = -1; \frac{1}{x} - \frac{2}{y} = 2, x, y \neq 0$$

OR

Find the value(s) of  $k$  for which the pair of equations  $\begin{cases} kx + 2y = 3 \\ 3x + 6y = 10 \end{cases}$  has a unique solution.

### SOLUTION:

The given equations are

$$\frac{3}{x} + \frac{8}{y} = -1 \quad \dots\dots (1)$$

$$\frac{1}{x} - \frac{2}{y} = 2 \quad \dots\dots (2)$$

Let  $\frac{1}{x} = u$  and  $\frac{1}{y} = v$

(1) and (2) will become

$$3u + 8v = -1 \quad \dots\dots (3)$$

$$u - 2v = 2 \quad \dots\dots (4)$$

Multiply (4) with 4

$$4u - 8v = 8 \quad \dots\dots (5)$$

Adding (3) and (5) we get

$$7u = 7$$

$$\Rightarrow u = 1$$

Putting this value in (4)

$$1 - 2v = 2$$

$$\Rightarrow v = \frac{-1}{2}$$

Now

$$\frac{1}{x} = u$$

$$\Rightarrow \frac{1}{x} = 1$$

$$\Rightarrow x = 1$$

And

$$\frac{1}{y} = v$$

$$\Rightarrow \frac{1}{y} = \frac{-1}{2}$$

$$\Rightarrow y = -2$$

OR

The given equations are

$$kx + 2y = 3$$

$$3x + 6y = 10$$

For a unique solution,

$$\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$$

where  $a_1 = k$ ,  $a_2 = 3$ ,  $b_1 = 2$ ,  $b_2 = 6$

$$\frac{k}{3} \neq \frac{2}{6}$$

$$\Rightarrow k \neq 1$$

For all values of  $k$  except 1, the given linear equations will have unique solution.

#### Question 10

How many multiples of 4 lie between 10 and 205 ?

OR

Determine the A.P. whose third term is 16 and 7<sup>th</sup> term exceeds the 5<sup>th</sup> by 12.

#### SOLUTION:

We need to find the number of multiples of 4 between 10 and 205.

So, multiples of 4 gives the sequence 12, 16, ..., 204

$$a = 12, d = 4 \text{ and } a_n = 204$$

$$\text{Using the formula } a_n = a + (n - 1)d$$

Plugging values in the formula we get

$$204 = 12 + (n - 1)4$$

$$204 = 12 + 4n - 4$$

$$4n = 196$$

$$n = 49$$

Thus, there are 49 multiples of 4 between 10 and 205.

OR

Given: 3rd term of the AP is 16.

$$a_3 = 16$$

$$a + (3 - 1)d = 16$$

$$a + 2d = 16 \quad \dots(1)$$

Also, 7th term exceeds the 5th term by 12.

$$a_7 - a_5 = 12$$

$$[a + (7 - 1)d] - [a + (5 - 1)d] = 12$$

$$(a + 6d) - (a + 4d) = 12$$

$$2d = 12$$

$$d = 6$$

From equation (1), we obtain

$$a + 2(6) = 16$$

$$a + 12 = 16$$

$$a = 4$$

Therefore, A.P. will be 4, 10, 16, 22, ...

### Question 11

Use Euclid's division algorithm to find the HCF of 255 and 867.

#### SOLUTION:

The given numbers are 255 and 867.

Now  $867 > 255$ . So, on applying Euclid's algorithm we get

$$867 = 255 \times 3 + 102$$

Now the remainder is not 0 so, we repeat the process again on 255 and 102

$$255 = 102 \times 2 + 51$$

The algorithm is applied again but this time on the numbers 102 and 51

$$102 = 51 \times 2 + 0$$

Thus, the HCF obtained is 51.

### Question 12

The point  $R$  divides the line segment  $AB$ , where  $A(-4, 0)$  and  $B(0, 6)$  such that  $AR = \frac{3}{4}AB$ . Find the coordinates of  $R$ .

**SOLUTION:**

We have given that R divides the line segment AB

$$AR + RB = AB$$

$$\frac{3}{4}AB + RB = AB$$

$$\Rightarrow RB = \frac{AB}{4}$$

$$\Rightarrow AR : RB = 3 : 1$$

Using section formula:

$$x = \left( \frac{m_1x_2 + m_2x_1}{m_1 + m_2} \right), y = \left( \frac{m_1y_2 + m_2y_1}{m_1 + m_2} \right)$$

$$m_1 = 3, m_2 = 1$$

$$x_1 = -4, y_1 = 0$$

$$x_2 = 0, y_2 = 6$$

Plugging values in the formula we get

$$x = \frac{3 \times 0 + 1 \times (-4)}{3 + 1}, y = \frac{3 \times 6 + 1 \times 0}{3 + 1}$$

$$x = \frac{-4}{4}, y = \frac{18}{4}$$

$$\Rightarrow x = -1, y = \frac{9}{2}$$

Therefore, the coordinates of R  $\left(-1, \frac{9}{2}\right)$ .

**Question 13**

Prove that:

$$(\sin \theta + 1 + \cos \theta) (\sin \theta - 1 + \cos \theta) \cdot \sec \theta \operatorname{cosec} \theta = 2$$

OR

Prove that :

$$\sqrt{\frac{\sec \theta - 1}{\sec \theta + 1}} + \sqrt{\frac{\sec \theta + 1}{\sec \theta - 1}} = 2 \operatorname{cosec} \theta$$

**SOLUTION:**

$$\text{LHS} = (\sin \theta + 1 + \cos \theta) (\sin \theta - 1 + \cos \theta) \cdot \sec \theta \operatorname{cosec} \theta$$

$$= [\sin^2 \theta - \sin \theta + \sin \theta \cos \theta + \sin \theta - 1 + \cos \theta + \sin \theta \cos \theta - \cos \theta + \cos^2 \theta] \frac{1}{\cos \theta} \frac{1}{\sin \theta} \quad \left( \because \sec \theta = \frac{1}{\cos \theta} \text{ and } \operatorname{cosec} \theta = \frac{1}{\sin \theta} \right)$$

$$= [1 + 2 \sin \theta \cos \theta - 1] \frac{1}{\cos \theta} \frac{1}{\sin \theta}$$

$$= [2 \sin \theta \cos \theta] \frac{1}{\cos \theta} \frac{1}{\sin \theta}$$

$$= 2 = \text{RHS}$$

Hence proved

OR



$$\begin{aligned}\sqrt{\frac{\sec \theta - 1}{\sec \theta + 1}} + \sqrt{\frac{\sec \theta + 1}{\sec \theta - 1}} &= \frac{\sqrt{\sec \theta - 1}}{\sqrt{\sec \theta + 1}} + \frac{\sqrt{\sec \theta + 1}}{\sqrt{\sec \theta - 1}} \\ &= \frac{\sqrt{\sec \theta - 1} \sqrt{\sec \theta - 1} + \sqrt{\sec \theta + 1} \sqrt{\sec \theta + 1}}{\sqrt{\sec \theta + 1} \sqrt{\sec \theta - 1}} \\ &= \frac{(\sqrt{\sec \theta - 1})^2 + (\sqrt{\sec \theta + 1})^2}{\sqrt{(\sec \theta - 1)(\sec \theta + 1)}}\end{aligned}$$

$$= \frac{\sec \theta - 1 + \sec \theta + 1}{\sqrt{\sec^2 \theta - 1}}$$

$$= \frac{2 \sec \theta}{\sqrt{\tan^2 \theta}}$$

$$= \frac{2 \sec \theta}{\tan \theta}$$

$$= 2 \frac{1}{\frac{\cos \theta}{\sin \theta}}$$

$$= 2 \frac{1}{\sin \theta}$$

$$= 2 \operatorname{cosec} \theta$$

**Question 14**

In what ratio does the point P(-4, y) divide the line segment joining the points A(-6, 10) and B(3, -8) ? Hence find the value of y.

**OR**

Find the value of p for which the points (-5, 1), (1, p) and (4, -2) are collinear.

**SOLUTION:**

Let P divides the line segment AB in the ratio  $k : 1$

Using section formula

$$x = \frac{m_1x_2+m_2x_1}{m_1+m_2}, y = \frac{m_1y_2+m_2y_1}{m_1+m_2}$$

A  $(-6, 10)$  and B  $(3, -8)$

$$m_1 : m_2 = k : 1$$

plugging values in the formula we get

$$-4 = \frac{k \times 3 + 1 \times (-6)}{k+1}, y = \frac{k \times (-8) + 1 \times 10}{k+1}$$

$$-4 = \frac{3k-6}{k+1}, y = \frac{-8k+10}{k+1}$$

Considering only  $x$  coordinate to find the value of  $k$

$$-4k - 4 = 3k - 6$$

$$-7k = -2$$

$$k = \frac{2}{7}$$


$$k : 1 = 2 : 7$$

Now, we have to find the value of  $y$

so, we will use section formula only in  $y$  coordinate to find the value of  $y$

$$y = \frac{2 \times (-8) + 7 \times 10}{2+7}$$

$$y = \frac{-16+70}{9}$$



$$y = 6$$

Therefore, P divides the line segment AB in 2 : 7 ratio  
And value of y is 6.

OR

Points are collinear means the area of triangle formed by the collinear points is 0.

Using

$$\begin{aligned} \text{area of triangle} &= \frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)] \\ &= \frac{1}{2} [-5(p - (-2)) + 1(-2 - 1) + 4(1 - p)] \\ &= \frac{1}{2} [-5(p + 2) + 1(-3) + 4(1 - p)] \\ &= \frac{1}{2} [-5p - 10 - 3 + 4 - 4p] \\ &= \frac{1}{2} [-5p - 9 - 4p] \end{aligned}$$

Area of triangle will be zero points being collinear

$$\frac{1}{2} [-5p - 4p - 9] = 0$$

$$\frac{1}{2} [-9p - 9] = 0$$

$$9p + 9 = 0$$


$$p = -1$$

Therefore, the value of  $p = -1$ .

#### Question 15

ABC is a right triangle in which  $\angle B = 90^\circ$ . If  $AB = 8$  cm and  $BC = 6$  cm, find the diameter of the circle inscribed in the triangle.

#### SOLUTION:



We have given that a circle is inscribed in a triangle

Using pythagoras theorem

$$(AC)^2 = (AB)^2 + (BC)^2$$

$$(AC)^2 = (8)^2 + (6)^2$$

$$(AC)^2 = 64 + 36$$

$$(AC)^2 = 100$$

$$\Rightarrow AC = 10$$

Area of  $\triangle ABC = \text{area of } \triangle APB + \text{area of } \triangle BPC + \text{area of } \triangle APC$

$$\frac{1}{2} \times b \times h = \frac{1}{2} \times b_1 \times h_1 + \frac{1}{2} \times b_2 \times h_2 + \frac{1}{2} \times b_3 \times h_3$$

$$\frac{1}{2} \times 6 \times 8 = \frac{1}{2} \times 8 \times r + \frac{1}{2} \times 6 \times r + \frac{1}{2} \times 10 \times r$$

$$24 = 4r + 3r + 5r$$

$$24 = 12r$$

$$\Rightarrow r = 2$$


$$\therefore d = 2r$$

$$\Rightarrow d = 2 \times 2$$

$$\Rightarrow d = 4 \text{ cm}$$

### Question 16

In Figure 1, BL and CM are medians of a  $\triangle ABC$  right-angled at A. Prove that  $4(BL^2 + CM^2) = 5BC^2$ .



OR

Prove that the sum of the squares of the sides of a rhombus is equal to the sum of the squares of its diagonals.

### SOLUTION:

To prove:  $4(BL^2 + CM^2) = 5BC^2$

Proof: In  $\triangle CAB$ ,

Applying Pythagoras theorem,

$$AB^2 + AC^2 = BC^2 \quad \dots(1)$$

In  $\triangle ABL$ ,

$$AL^2 + AB^2 = BL^2$$

$$\Rightarrow \left(\frac{AC}{2}\right)^2 + AB^2 = BL^2$$

$$\Rightarrow AC^2 + 4AB^2 = 4BL^2 \quad \dots(2)$$

In  $\triangle CAM$ ,

$$CA^2 + MA^2 = CM^2$$

$$\Rightarrow \left(\frac{BA}{2}\right)^2 + CA^2 = CM^2$$

$$\Rightarrow BA^2 + 4CA^2 = 4CM^2 \quad \dots\dots(3)$$

Adding (2) and (3)


$$AC^2 + 4AB^2 + BA^2 + 4CA^2 = 4BL^2 + 4CM^2$$

$$\Rightarrow 5AC^2 + 5AB^2 = 4(BL^2 + CM^2)$$

$$\Rightarrow 5(AC^2 + AB^2) = 4(BL^2 + CM^2)$$

$$\Rightarrow 5(BC^2) = 4(BL^2 + CM^2) \quad \text{(From (1))}$$

Hence Proved.



In  $\triangle AOB$ ,  $\triangle BOC$ ,  $\triangle COD$ ,  $\triangle AOD$ ,

Applying Pythagoras theorem, we obtain

$$AB^2 = AO^2 + OB^2 \quad \dots (1)$$

$$BC^2 = BO^2 + OC^2 \quad \dots (2)$$

$$CD^2 = CO^2 + OD^2 \quad \dots (3)$$

$$AD^2 = AO^2 + OD^2 \quad \dots (4)$$

Adding all these equations, we obtain

$$AB^2 + BC^2 + CD^2 + AD^2 = 2(AO^2 + OB^2 + OC^2 + OD^2)$$

$$= 2\left(\left(\frac{AC}{2}\right)^2 + \left(\frac{BD}{2}\right)^2 + \left(\frac{AC}{2}\right)^2 + \left(\frac{BD}{2}\right)^2\right)$$


(Diagonals bisect each other)

$$= 2\left(\frac{(AC)^2}{2} + \frac{(BD)^2}{2}\right)$$


$$= (AC)^2 + (BD)^2$$

**Question 17**

In Figure 2, two concentric circles with centre O, have radii 21 cm and 42 cm. If  $\angle AOB = 60^\circ$ , find the area of the shaded region.



**SOLUTION:**



Radius of inner circle,  $OC = 21$  cm

Radius of outer circle,  $OA = 42$  cm

Area of circle with radius  $R = \pi R^2 = \pi(42)^2$

Area of circle with radius  $r = \pi r^2 = \pi(21)^2$

Area of sector AOB =  $\frac{\theta}{360} \times \pi R^2 = \frac{60}{360} \times \pi(42)^2 = \frac{\pi(42)^2}{6}$

Area of sector COD =  $\frac{\theta}{360} \times \pi r^2 = \frac{60}{360} \times \pi(21)^2 = \frac{\pi(21)^2}{6}$

Area of shaded portion = Area of circle with radius  $R$  – Area of circle with radius  $r$  – [Area of sector AOB – Area of sector COD]

$$\begin{aligned}
 &= \pi(42)^2 - \pi(21)^2 - \left[ \frac{\pi(42)^2}{6} - \frac{\pi(21)^2}{6} \right] \\
 &= \pi \left[ (42)^2 - (21)^2 - \frac{1}{6} \left[ (42)^2 - (21)^2 \right] \right] \\
 &= \pi \left[ \left( (42)^2 - (21)^2 \right) \left( 1 - \frac{1}{6} \right) \right] \\
 &= \pi \left[ (42 - 21)(42 + 21) \frac{5}{6} \right] \\
 &= \frac{22}{7} \times \frac{5}{6} \times 21 \times 63 \\
 &= 3465 \text{ cm}^2
 \end{aligned}$$

Question 18

Calculate the mode of the following distribution :

Class :	10 - 15	15 - 20	20 - 25	25 - 30	30 - 35
Frequency :	4	7	20	8	1

SOLUTION:

Modal class is the class with highest frequency

modal class is 20 - 25

lower limit of modal class i.e  $l = 20$

class size i.e  $h = 5$

frequency of modal class  $f_1 = 20$

frequency of preceding class  $f_0 = 7$

frequency of succeeding class  $f_2 = 8$

Using the formula

$$\text{mode} = l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h$$

Plugging the values in the formula we get

$$\text{mode} = 20 + \left( \frac{20 - 7}{2 \times 20 - 7 - 8} \right) \times 5$$

$$\text{mode} = 20 + \left( \frac{13}{25} \right) \times 5$$

$$\text{mode} = 20 + \frac{13}{5}$$

$$\text{mode} = \frac{113}{5} = 22.6$$

Question 19

A cone of height 24 cm and radius of base 6 cm is made up of modelling clay. A child reshapes it in the form of a sphere. Find the radius of the sphere and hence find the surface area of this sphere.

OR

A farmer connects a pipe of internal diameter 20 cm from a canal into a cylindrical tank

in his field which is 10 m in diameter and 2 m deep. If water flows through the pipe at the rate of 3 km/hr, how much time will the tank be filled ?

**SOLUTION:**

A cone has been reshaped in sphere  
Height of cone is 24 cm and radius of base is 6 cm  
Volume of sphere = volume of cone  
Volume of cone =  $\frac{1}{3}\pi r^2 h$   
Plugging the values in the formula we get  
volume of cone =  $\frac{1}{3}\pi(6)^2 24$

$$= 288\pi \text{ cm}^3$$

Let the radius of sphere be  $r$

$$\text{Volume of sphere} = \frac{4}{3}\pi r^3$$

Since, volume of cone = volume of sphere

$$\text{Volume of sphere} = 288\pi \text{ cm}^3$$

So,

$$288\pi = \frac{4}{3}\pi r^3$$

$$\Rightarrow 288 = \frac{4}{3}r^3$$

$$\Rightarrow r^3 = 216$$

$$\Rightarrow r = 6 \text{ cm}$$

Hence, radius of reshaped sphere is 6 cm

$$\text{Now, surface area of sphere} = 4\pi r^2$$

$$= 4\pi(6)^2$$

$$= 144 \times \frac{22}{7}$$

$$= 452.5 \text{ cm}^2$$

Therefore, surface area of sphere is  $452.57 \text{ cm}^2$ .

OR



Consider an area of cross-section of pipe as shown in the figure.

$$\text{Radius } (r_1) \text{ of circular end of pipe} = \frac{20}{200} = 0.1 \text{ m}$$


$$\text{Area of cross-section} = \pi \times r_1^2 = \pi \times (0.1)^2 = 0.01\pi \text{ m}^2$$



$$\text{Speed of water} = 3 \text{ km/h} = \frac{3000}{60} = 50 \text{ metre/min}$$

$$\text{Volume of water that flows in 1 minute from pipe} = 50 \times 0.01\pi = 0.5\pi \text{ m}^3$$

$$\text{Volume of water that flows in } t \text{ minutes from pipe} = t \times 0.5\pi \text{ m}^3$$



$$\text{Radius } (r_2) \text{ of circular end of cylindrical tank} = \frac{10}{2} = 5 \text{ m}$$

$$\text{Depth } (h_2) \text{ of cylindrical tank} = 2 \text{ m}$$

Let the tank be filled completely in  $t$  minutes.

Volume of water filled in tank in  $t$  minutes is equal to the volume of water flowed in  $t$  minutes from the pipe.

$$\text{Volume of water that flows in } t \text{ minutes from pipe} = \text{Volume of water in tank}$$

$$t \times 0.5\pi = \pi \times (r_2)^2 \times h_2$$

$$t \times 0.5 = 5^2 \times 2$$

$$t = 100$$

Therefore, the cylindrical tank will be filled in 100 minutes.

Question 20

Prove that  $2 + 3\sqrt{3}$  is an irrational number when it is given that  $\sqrt{3}$  is an irrational number.

SOLUTION:

To prove:  $2 + 3\sqrt{3}$  is irrational, let us assume that  $2 + 3\sqrt{3}$  is rational.

$2 + 3\sqrt{3} = \frac{a}{b}$ ;  $b \neq 0$  and  $a$  and  $b$  are integers.

$$\Rightarrow 2b + 3\sqrt{3}b = a$$

$$\Rightarrow 3\sqrt{3}b = a - 2b$$

$$\Rightarrow \sqrt{3} = \frac{a-2b}{3b}$$

Since  $a$  and  $b$  are integers so,  $a - 2b$  will also be an integer.

So,  $\frac{a-2b}{3b}$  will be rational which means  $\sqrt{3}$  is also rational.

But we know  $\sqrt{3}$  is irrational(given).

Thus, a contradiction has risen because of incorrect assumption.

Thus,  $2 + 3\sqrt{3}$  is irrational.

Question 21

Sum of the areas of two squares is  $157 \text{ m}^2$ . If the sum of their perimeters is 68 m, find the sides of the two squares.

SOLUTION:

Let the side of one square be  $x$

And side of other square be  $y$

Sum of area of two square is 157

Equation becomes

$$x^2 + y^2 = 157 \quad \dots\dots(1) \quad (\because \text{area of square is side}^2)$$

Now, sum of their perimeters is 68

Equation becomes

$$4x + 4y = 68 \quad (\because \text{perimeter of square is } 4 \times \text{side})$$

solving the two equation by substitution method

$$4x + 4y = 68$$

$$x + y = 17$$

$$\Rightarrow x = 17 - y \quad \dots\dots(2)$$

Substitute (2) in (1)

$$(17 - y)^2 + y^2 = 157$$

$$289 + y^2 - 34y + y^2 = 157$$

$$2y^2 - 34y + 132 = 0$$

$$y^2 - 17y + 66 = 0$$

$$\text{Using } y = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Plugging the values in the formula we get

$$y = \frac{17 \pm \sqrt{289 - 4(66)}}{2}$$

$$y = \frac{17 \pm \sqrt{25}}{2}$$

$$y = \frac{17 \pm 5}{2}$$

$$y = \frac{12}{2}, \frac{22}{2}$$

$$y = 6, 11$$

when  $y = 6$  then  $x = 11$

And when  $y = 11$  then  $x = 6$

Therefore, the sides of square are 6 m and 11 m.

### Question 22

Find the quadratic polynomial, sum and product of whose zeroes are  $-1$  and  $-20$  respectively. Also find the zeroes of the polynomial so obtained.

### SOLUTION:

We have been given the sum of zeroes and product of zeroes

Let us consider the general polynomial

$$p(x) = ax^2 + bx + c$$

Sum of zeroes is  $-\frac{b}{a}$

And product of zeroes is  $\frac{c}{a}$

According to question

$$\frac{-b}{a} = -1 \text{ and } \frac{c}{a} = -20$$

Assuming  $a = 1$

$$-b = -1$$

$$\Rightarrow b = 1$$

And  $c = -20$

So, the polynomial so formed is  $p(x) = x^2 + x - 20$

To find the zeroes of the polynomial equate polynomial to zero.

$$x^2 + x - 20 = 0$$

$$x^2 + 5x - 4x - 20 = 0$$

$$x(x + 5) - 4(x + 5) = 0$$

$$(x + 5)(x - 4) = 0$$

$$\Rightarrow x = -5, 4$$

Therefore, zeroes of the polynomial are  $-5$  and  $4$ .

Question 23

A plane left 30 minutes later than the scheduled time and in order to reach its destination 1500 km away on time, it has to increase its speed by 250 km/hr from its usual speed. Find the usual speed of the plane.

OR

Find the dimensions of a rectangular park whose perimeter is 60 m and area 200 m<sup>2</sup>.

SOLUTION:

Let the usual speed of the plane be  $x$  km/hr

And the new speed of the plane after increased by 250 is  $(x + 250)$  km / hr

According to question

$$\frac{1500}{x} - \frac{1500}{(x+250)} = \frac{30}{60}$$

$$\Rightarrow \frac{1500x + 1500 \times 250 - 1500x}{x(x+250)} = \frac{1}{2}$$

$$\Rightarrow 1500 \times 250 \times 2 = x(x + 250)$$

$$\Rightarrow 750000 = x^2 + 250x$$

$$\Rightarrow x^2 + 1000x - 750x - 750000 = 0$$

$$\Rightarrow (x + 1000)(x - 750) = 0$$

$$x = 750, -1000$$

Speed can not be negative so -1000 will be neglected

Therefore, usual speed of the plane is 750 km/hr.

OR

Let the length of rectangle be  $x$

And breadth of rectangle be  $y$

$$xy = 200 \quad (\because \text{area} = \text{length} \times \text{breadth})$$

$$\text{And } 2(x + y) = 60 \quad [\because \text{perimeter} = 2(\text{length} + \text{breadth})]$$

$$\text{substitute } y = \frac{200}{x} \text{ in } 2(x + y) = 60$$

Equation becomes:

$$2\left(x + \frac{200}{x}\right) = 60$$

$$2\left(\frac{x^2 + 200}{x}\right) = 60$$

$$2x^2 - 60x + 400 = 0$$

$$x^2 - 30x + 200 = 0$$

Using

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Plugging the values we get:

$$x = \frac{30 \pm \sqrt{(-30)^2 - 4(1)(200)}}{2}$$

$$x = \frac{30 \pm 10}{2}$$

$$x = \frac{40}{2}, \frac{20}{2}$$

$$x = 20, 10$$

when  $x = 20$  then  $y = 10$

And when  $x = 10$  then  $y = 20$ .

### Question 24

Find the value of  $x$ , when in the A.P. given below  
 $2 + 6 + 10 + \dots + x = 1800$ .

### SOLUTION:

We have been given an A.P

$$2 + 6 + 10 + \dots + x = 1800$$

$$a = 2, d = 6 - 2 = 4, a_n = x \text{ and } s_n = 1800$$

Firstly, we will find using

$$S_n = \frac{n}{2} [2a + (n - 1)d]$$

$$1800 = \frac{n}{2} [2 \times 2 + (n - 1)4]$$

$$1800 = \frac{4n + 4n^2 - 4n}{2}$$

$$900 = n^2$$

$$\Rightarrow n = \pm 30$$

Number of terms can not be negative

$$n = 30$$

Now for value of  $x$  which is  $a_n$

$$a_n = a + (n - 1)d$$

$$x = 2 + (30 - 1)4$$

$$x = 2 + 116$$

$$x = 118$$

Therefore, value of  $x$  is 118.

Question 25

If  $\sec \theta + \tan \theta = m$ , show that  $\frac{m^2-1}{m^2+1} = \sin \theta$ .


SOLUTION:

$$\begin{aligned} & \frac{m^2-1}{m^2+1} \\ \Rightarrow & \frac{(\sec \theta + \tan \theta)^2 - (\sec^2 \theta - \tan^2 \theta)}{(\sec \theta + \tan \theta)^2 + (\sec^2 \theta - \tan^2 \theta)} \\ \Rightarrow & \frac{\sec^2 \theta + \tan^2 \theta + 2 \sec \theta \tan \theta - \sec^2 \theta + \tan^2 \theta}{\sec^2 \theta + \tan^2 \theta + 2 \sec \theta \tan \theta + \sec^2 \theta - \tan^2 \theta} \\ \Rightarrow & \frac{2 \tan \theta (\tan \theta + \sec \theta)}{2 \sec \theta (\tan \theta + \sec \theta)} \\ \Rightarrow & \frac{\tan \theta}{\sec \theta} = \frac{\sin \theta}{\cos \theta \sec \theta} \quad (\because \tan \theta = \frac{\sin \theta}{\cos \theta}) \\ \Rightarrow & \frac{\sin \theta}{\cos \theta \times \frac{1}{\cos \theta}} = \sin \theta \quad (\because \cos \theta = \frac{1}{\sec \theta}) \end{aligned}$$

Hence, proved

Question 26

In  $\triangle ABC$  (Figure 3),  $AD \perp BC$ . Prove that  $AC^2 = AB^2 + BC^2 - 2BC \times BD$



SOLUTION:

Applying Pythagoras theorem in  $\triangle ADB$ , we obtain

$$AD^2 + DB^2 = AB^2$$

$$\Rightarrow AD^2 = AB^2 - DB^2 \quad \dots(1)$$

Applying Pythagoras theorem in  $\triangle ADC$ , we obtain

$$AD^2 + DC^2 = AC^2$$

$$AB^2 - BD^2 + DC^2 = AC^2 \text{ [Using equation (1)]}$$

$$AB^2 - BD^2 + (BC - BD)^2 = AC^2$$

$$AC^2 = AB^2 - BD^2 + BC^2 + BD^2 - 2BC \times BD$$

$$AC^2 = AB^2 + BC^2 - 2BC \times BD$$


**Question 27**

A moving boat is observed from the top of a 150 m high cliff moving away from the cliff. The angle of depression of the boat changes from  $60^\circ$  to  $45^\circ$  in 2 minutes. Find the speed of the boat in m/min.

OR

There are two poles, one each on either bank of a river just opposite to each other. One pole is 60 m high. From the top of this pole, the angle of depression of the top and foot of the other pole are  $30^\circ$  and  $60^\circ$  respectively. Find the width of the river and height of the other pole.

**SOLUTION:**



Let AO be the cliff of height 150 m.

Let the speed of boat be  $x$  metres per minute.

And BC be the distance which man travelled.

So,  $BC = 2x$  [ $\because$  Distance = Speed  $\times$  Time]

$$\tan(60^\circ) = \frac{AO}{OB}$$

$$\sqrt{3} = \frac{150}{OB}$$

$$\Rightarrow OB = \frac{150\sqrt{3}}{3} = 50\sqrt{3}$$

$$\tan(45^\circ) = \frac{AO}{OC}$$

$$\Rightarrow 1 = \frac{150}{OC}$$

$$\Rightarrow OC = 150$$

Now  $OC = OB + BC$

$$\Rightarrow 150 = 50\sqrt{3} + 2x$$

$$\Rightarrow x = \frac{150 - 50\sqrt{3}}{2}$$


$$\Rightarrow x = 75 - 25\sqrt{3}$$

Using  $\sqrt{3} = 1.73$

$$x = 75 - 25 \times 1.732 \approx 32 \text{ m/min}$$

Hence, the speed of the boat is 32 metres per minute.

OR



Let the width of the river be  $w$ .

In  $\triangle ABC$ ,

$$\tan 60^\circ = \frac{AB}{BC}$$

$$\Rightarrow \sqrt{3} = \frac{60}{w}$$

$$\Rightarrow w = \frac{60}{\sqrt{3}} = \frac{60\sqrt{3}}{3} = 20\sqrt{3}$$

In  $\triangle AED$ ,

$$\tan 30^\circ = \frac{AE}{ED}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{AE}{w}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{AE}{20\sqrt{3}}$$



$$\Rightarrow AE = 20$$

Height of pole  $CD = AB - AE$

$$= 60 - 20 = 40 \text{ m}$$


Thus, width of river is  $20\sqrt{3} = 20 \times 1.732 = 34.64 \text{ m}$

Height of pole = 40 m

### Question 28

Construct a triangle with sides 5 cm, 6 cm and 7 cm and then another triangle whose sides are  $\frac{3}{5}$  of the corresponding sides of the first triangle.

### SOLUTION:



1. Draw a line  $AB = 5 \text{ cm}$  and draw a ray from A and taking A as centre cut an arc at C of 6 cm and taking B as centre cut an arc of 7 cm at C
2. Draw  $AX$  such that  $\angle BAX$  is an acute angle.
3. Cut 5 equal arcs  $AA_1, A_1A_2, A_2A_3, A_3A_4$  and  $A_4A_5$ .
4. Join  $A_5$  to B and draw a line through  $A_3$  parallel to  $A_5B$  which meets AB at  $B'$ .

$$\text{Here, } AB' = \frac{3}{5} AB$$

5. Now draw a line through  $B'$  parallel to  $BC$  which joins  $AC$  at  $C'$ .

$$\text{Here, } B'C' = \frac{3}{5} BC \text{ and } AC' = \frac{3}{5} AC$$

Thus,  $AB'C'$  is the required triangle.

### Question 29

## QB365-Question Bank Software

Calculate the mean of the following frequency distribution:

Class:	10-30	30-50	50-70	70-90	90-110	110-130
Frequency:	5	8	12	20	3	2

**OR**

The following table gives production yield in kg per hectare of wheat of 100 farms of a village:

Production yield (kg/hectare):	40-45	45-50	50-55	55-60	60-65	65-70
Number of farms	4	6	16	20	30	24

Change the distribution to a 'more than type' distribution, and draw its ogive.

**SOLUTION:**

Class	frequency ( $f_i$ )	Class mark ( $x_i$ )	$f_i x_i$
10-30	5	$\frac{10+30}{2} = 20$	100
30-50	8	$\frac{30+50}{2} = 40$	320
50-70	12	$\frac{50+70}{2} = 60$	720
70-90	20	$\frac{70+90}{2} = 80$	1600
90-110	3	$\frac{90+110}{2} = 100$	300
110-130	2	$\frac{110+130}{2} = 120$	240
	$\sum f_i = 50$		$\sum f_i x_i = 3280$


Using: mean =  $\frac{\sum f_i x_i}{\sum f_i}$

substituting the values in the formula

mean =  $\frac{3280}{50} = 65.6$

**OR**

Production yield	Cumulative frequency
more than 40	100
more than 45	96
more than 50	90
more than 55	74
more than 60	54
more than 65	24



**Question 30**

A container opened at the top and made up of a metal sheet, is in the form of a frustum of a cone of height 16 cm with radii of its lower and upper ends as 8 cm and 20 cm respectively. Find the cost of milk which can completely fill the container, at the rate of ₹ 50 per litre. Also find the cost of metal sheet used to make the container, if it costs ₹ 10 per 100 cm<sup>2</sup>. (Take  $\pi = 3.14$ )

**SOLUTION:**

We have to find the cost of milk which can completely fill the container

Volume of container = Volume of frustum

$$= \frac{1}{3}\pi h (r_1^2 + r_2^2 + r_1 r_2)$$

Here,

height = 16 cm

radius of upper end = 20 cm

And radius of lower end = 8 cm

Plugging the values in the formula we get

$$\begin{aligned}\text{Volume of container} &= \frac{1}{3} \times 3.14 \times 16 \left( (20)^2 + (8)^2 + 20 \times 8 \right) \\ &= \frac{1}{3} \times 50.24 (400 + 64 + 160) \\ &= \frac{1}{3} \times 50.24 (624) \\ &= 10449.92 \text{ cm}^3 \\ &= 10.449 \text{ litre} \quad (\because 1 \text{ litre} = 1000 \text{ cm}^3)\end{aligned}$$

Cost of 1 litre milk is Rs 50

Cost of 10.449 litre milk =  $50 \times 10.449 = \text{Rs } 522.45$

We will find the cost of metal sheet to make the container

Firstly, we will find the area of container

Area of container = Curved surface area of the frustum + area of bottom circle ( $\because$  container is closed from bottom)

$$\text{Area of container} = \pi (r_1 + r_2)l + \pi r^2$$

Now, we will find  $l$

$$l = \sqrt{h^2 + (r_1 - r_2)^2}$$

$$l = \sqrt{(16)^2 + (20 - 8)^2}$$

$$l = \sqrt{(16)^2 + (12)^2}$$

$$l = \sqrt{256 + 144}$$

$$l = \sqrt{400}$$

$$l = 20 \text{ cm}$$

$$\begin{aligned}\text{Area of frustum} &= 3.14 \times 20 (20 + 8) \\ &= 1758.4 \text{ cm}^2\end{aligned}$$

$$\text{Area of bottom circle} = 3.14 \times 8^2 = 200.96 \text{ cm}^2$$

$$\begin{aligned}\text{Area of container} &= 1758.4 + 200.96 \\ &= 1959.36 \text{ cm}^2\end{aligned}$$

$$\text{Cost of making } 100 \text{ cm}^2 = \text{Rs } 10$$

$$\text{Cost of making } 1 \text{ cm}^2 = \frac{10}{100} = \text{Rs } \frac{1}{10}$$

$$\text{Cost of making } 1959.36 \text{ cm}^2 = \frac{1}{10} \times 1959.36 = 195.936$$

Hence, cost of milk is Rs 522.45

And cost of metal sheet is Rs 195.936