

QB365 - Question Bank Software

CLASS XII SESSION 2020-21

PRACTICE PAPER 5


SUBJECT : MATHEMATICS

MARKING SCHEME (THEORY)

Sr No	Objective type Question	Marks
	Section I	
1	$2^3 A = 8 \times 5$ OR $x=y$	1
2	$A_1 \cup A_2 \cup A_3 = A$ and $A_1 \cap A_2 \cap A_3 = \emptyset$	1
3	$A=2, b=3$ OR $1-\alpha^2-\beta\gamma=0$	1
4	Reflexive Relation	1
5	R is not symmetric	1
6	$3 \times p$	1
7	$\frac{1}{3}e^{x^3} + C$ OR $e-1$	1
8	$\frac{1}{4}$	1
9	1	1
10	$\vec{r} = (3\hat{i} + 4\hat{j} + 5\hat{k}) + \lambda(2\hat{i} + 2\hat{j} - 3\hat{k})$ OR distance = $\sqrt{b^2 + c^2}$	1
11	Order = 2, Degree 1	1
12	-1	1
13	P = -2	1
14	Zero OR 3	1
15	$\frac{2}{3}$ Square Units OR $\frac{37}{3}$ Units	1
16	$n=3$ OR $\log x$	1
17	CASE STUDY - I	1
	(i) $2x+2y=36 \Rightarrow x+y=18 \Rightarrow y=18-x$	1
	(ii) $V(x) = \pi x^2 y = \pi x^2 [18-x] = \pi [18x^2 - x^3]$	1
	(iii) $V(x) = \pi(36x - 3x^2) = 0 \Rightarrow 36x = 3x^2 \Rightarrow x=0, x=12$ $X=12, y=6$	1
	(iv) $V(x) = \pi [18 \cdot 12^2 - 12^3]$ $= \pi 12^2 [18 - 12]$ $= \pi \cdot 144 \cdot 6 = 864\pi$	1
	(v) $\frac{dv}{dx} = \pi(36x - 3x^2)$ $\frac{d^2v}{dx^2} = \pi(36 - 6x) = -36\pi < 0$	1
18	CASE STUDY - II	1
	(i) $P(U_1) = P(U_2) = P(U_3) = \frac{1}{3}$ Equal probability	1
	(ii) E_1 = Event of a ball chosen from $P(E_1) = P(E_2) = P(E_3) = \frac{1}{3}$	1
	(III) $P\left(\frac{E}{E_1}\right) = \frac{2}{5}, P\left(\frac{E}{E_2}\right) = \frac{3}{5}, P\left(\frac{E}{E_3}\right) = \frac{4}{5}$	1

QB365 - Question Bank Software

	<p>E-White ball</p> <p>(iv) $P\left(\frac{E_2}{E_1}\right) = \frac{P(E_2) P\left(\frac{E_2}{E_1}\right)}{P(E_1) \cdot P\left(\frac{E_2}{E_1}\right) + P(E_2) \cdot P\left(\frac{E_2}{E_2}\right) + P(E_3) \cdot P\left(\frac{E_2}{E_3}\right)}$</p> $= \frac{\frac{1}{3} \cdot \frac{3}{5}}{\frac{1}{3} \cdot \frac{3}{5} + \frac{1}{3} \cdot \frac{1}{3} + \frac{1}{3} \cdot \frac{1}{5}} = \frac{\frac{1}{5}}{\frac{1}{5} + \frac{1}{9} + \frac{1}{15}} = \frac{3}{9} = \frac{1}{3}$ <p>(v) $P(E_2) = 1/3$ $P(E/E_2) = 3/5$ $P(E_2) P(E/E_2) = 1/3 \times 3/5 = 1/5$</p>	<p>1</p> <p>1</p> <p>1</p>
	<p>PART B</p> <p>SECTION III 2 M</p>	
19	$\tan^{-1} \left[\frac{\cos x}{1 - \sin x} \right] = \tan^{-1} \left[\frac{\sin(\frac{\pi}{2} - x)}{1 - \cos(\frac{\pi}{2} - x)} \right] = \tan^{-1} \left[\frac{2 \sin(\frac{\pi}{4} - \frac{x}{2}) \cos(\frac{\pi}{4} - \frac{x}{2})}{2 \sin^2(\frac{\pi}{4} - \frac{x}{2})} \right]$ $= \tan^{-1} \left[\cot\left(\frac{\pi}{4} - \frac{x}{2}\right) \right] = \tan^{-1} \left[\tan\left(\frac{\pi}{2} - \left(\frac{\pi}{4} - \frac{x}{2}\right)\right) \right]$ $= \frac{\pi}{4} + \frac{x}{2}$	<p>1</p> <p>1</p>
20	<p>2A-3B+5C=0</p> $\Rightarrow 2A=3B-5C=3 \begin{bmatrix} -2 & 2 & 0 \\ 3 & 1 & 4 \end{bmatrix} - 5 \begin{bmatrix} 2 & 0 & -2 \\ 7 & 1 & 6 \end{bmatrix}$ $\Rightarrow 2A = \begin{bmatrix} -16 & 6 & 10 \\ -26 & -2 & -18 \end{bmatrix} \Rightarrow A = \begin{bmatrix} -8 & 3 & 5 \\ -13 & -1 & -9 \end{bmatrix}$ <p align="center">OR</p> $\begin{bmatrix} 2x \\ x \end{bmatrix} + \begin{bmatrix} 3y \\ 5y \end{bmatrix} + \begin{bmatrix} -8 \\ -11 \end{bmatrix} = 0$ <p>2x+3y-8=0 X+5y-11=0 solving x=1, y=2</p>	<p>½</p> <p>1</p> <p>½</p> <p>½</p> <p>1</p> <p>1/2</p>
21	K=6	2
22	$\frac{dy}{dx} = \frac{-16x}{9y}$ slope of tangent at(2,3)=-32/27 , Slope of normal=27/32 Equation of tangent=32x+27y=145 , Equation of normal=27x-32y=-4	1+1
23	$\int \frac{1-2\sin^2 x + 2\sin^4 x}{\cos^2 x} dx = \int \sec^2 x dx = \tan x + C$ <p align="center">OR</p> $\frac{1}{4} \int_0^a \frac{dx}{\frac{1}{4} + x^2} = \frac{\pi}{8}$ $\Rightarrow \frac{1}{4} \int_0^a \frac{dx}{\left(\frac{1}{2}\right)^2 + x^2} = \frac{\pi}{8}$ $\Rightarrow \frac{1}{4} \left[\frac{a \tan^{-1} 2x}{1/2} \right] = \frac{\pi}{8} \Rightarrow \frac{1}{4} [\tan^{-1} 2a - \tan^{-1} 0] = \frac{\pi}{8} \Rightarrow a = 1/2$	<p>2</p> <p>½</p> <p>1+1/2</p>


24	<p>We have $y^2 = x$ and $2y = x$</p> <p>Solving, we get $y^2 = 2y$ $\Rightarrow y = 0, 2$ When $y = 0, x=0$ and when $y = 2, x = 4$ So, points of intersection are $(0,0)$ and $(4,2)$ Graphs of parabola $y^2 = x$ and $2y = x$ are as shown in the following figure.</p>  <p>From the figure, area of the shaded region,</p> $A = \int_0^4 \left[\sqrt{x} - \frac{x}{2} \right] dx$ $= \left[\frac{2}{3} x^{3/2} - \frac{1}{2} \cdot \frac{x^2}{2} \right]_0^4 = \frac{2}{3} \cdot 4^{3/2} - \frac{16}{4} - 0 = \frac{16}{3} - 4 = \frac{4}{3} \text{ sq. units}$	1+1
25	$\frac{dy}{dx} + 2y \tan x = \sin x \quad P=2 \tan x \quad Q=\sin x$ $IF = e^{\int 2 \tan x dx} = e^{2 \int \tan x dx} = e^{2 \log \sec x} = \sec^2 x$ <p>Solution $= y \sec^2 x = \int \sin x \sec^2 x dx = \int \sec x \tan x dx \Rightarrow y \sec^2 x = \sec x + C$ at $x = \frac{\pi}{3}, y=0 \Rightarrow 0 = \sec \frac{\pi}{3} + C \Rightarrow C = -2$ PS $= y \sec^2 x = \sec x - 2$</p>	1+1
26	$\vec{PQ} = \hat{i} - 4\hat{j} - 2\hat{k}$ $\vec{PR} = -3\hat{i} + 12\hat{j} + 6\hat{k} = -3[\hat{i} - 4\hat{j} - 2\hat{k}]$ $\Rightarrow \vec{PR} = -3\vec{PQ}$ $\Rightarrow P, Q, R \text{ are collinear}$	1+1
27	<p>Normal vector $= \vec{n}$ Through $(1,0,0)$ ie \hat{i}</p> $(\vec{r} - \hat{i}) \cdot \vec{n} = 0 \quad \text{Plain contains line } \vec{r} = 0 + \lambda \hat{j}$ $(\vec{r} - \hat{i}) \cdot \hat{k} = 0 \quad i.n=0 \quad \vec{n} = \hat{k}$ $r.\hat{k} = 0 \quad j.n=0$	1+1

QB365 - Question Bank Software

28	$P\left(\frac{\bar{E}}{\bar{F}}\right) = \frac{P(\bar{E} \cap \bar{F})}{P(\bar{F})} = \frac{P(E \cup F)}{P(\bar{F})} = \frac{1 - P(E \cap F)}{1 - P(F)}$ $P(E \cup F) = P(E) + P(F) - P(E \cap F)$ $= 0.8 + 0.7 - 0.6 = 0.9$ $\Rightarrow P\left(\frac{\bar{E}}{\bar{F}}\right) = \frac{1 - 0.9}{1 - 0.7} = \frac{0.1}{0.3} = \frac{1}{3}$ <p style="text-align: center;">OR</p> $P(A \cap B) = \frac{7}{10}, P(B) = \frac{17}{20}$ $P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)}$ $= \frac{\frac{7}{10}}{\frac{17}{20}} = \frac{14}{17}$	1+1 1 1
	SECTION IV	
29	<p style="text-align: center;">For $x_1, x_2 \in A$</p> <p>(i) $f(x_1) = f(x_2) \Rightarrow \frac{4x_1+3}{6x_1-4} = \frac{4x_2+3}{6x_2-4}$</p> $\Rightarrow 34x_1 = 34x_2$ $\Rightarrow x_1 = x_2 \quad \text{f is one-one}$ <p>(ii) $y = \frac{4x+3}{6x-4} \Rightarrow x = \frac{4y+3}{6y-4}$ f is onto</p>	1+1+1
30	<p style="text-align: center;">Point of intersection $= x = y^2 \quad xy = k$</p> <p>(1) $y^2 y = k \Rightarrow y = k^{\frac{1}{3}}, x = k^{\frac{2}{3}}$</p> <p>(2) $m_1 = \frac{1}{2y}, m_2 = \frac{-y}{x}$</p> <p>(3) $m_1 m_2 = -1 = \frac{1}{2y} \left(\frac{-y}{x}\right) = -1 \Rightarrow 2x = 1$</p> <p>$2k^{\frac{2}{3}} = 1$ cubing $8k^2 = 1$</p> <p>OR</p> $y = \tan^{-1} \sqrt{\frac{1 - \cos x}{1 + \cos x}} \Rightarrow y = \tan^{-1} \sqrt{\frac{\sin^2 \frac{x}{2}}{\cos^2 \frac{x}{2}}}$ $\Rightarrow y = \tan^{-1} \left(\tan \frac{x}{2} \right)$ $\Rightarrow y = \frac{x}{2} \Rightarrow \frac{dy}{dx} = \frac{1}{2}$	1+1+1 1 1 1

31

1+1+1



$$\therefore \text{Required Area} = \int_0^4 x \, dx + \int_4^{4\sqrt{2}} \sqrt{(4\sqrt{2})^2 - x^2} \, dx \quad \dots(1)$$

Taking I_1 i.e.

$$I_1 = \int_0^4 x \, dx$$

$$= \left[\frac{x^2}{2} \right]_0^4$$

$$= \frac{(4)^2 - 0}{2}$$

$$= \frac{16}{2}$$

$$= 8$$

Now solving I_2


$$I_2 = \int_4^{4\sqrt{2}} \sqrt{(4\sqrt{2})^2 - x^2} dx$$

It is of form

$$\int \sqrt{a^2 - x^2} dx = \frac{1}{2} x \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + c$$

Replacing a by $4\sqrt{2}$, we get

$$\begin{aligned} I_2 &= \left[\frac{x}{2} \sqrt{(4\sqrt{2})^2 - x^2} + \frac{(4\sqrt{2})^2}{2} \sin^{-1} \frac{x}{4\sqrt{2}} \right]_4^{4\sqrt{2}} \\ &= \frac{4\sqrt{2}}{2} \sqrt{(4\sqrt{2})^2 - (4\sqrt{2})^2} + \frac{(4\sqrt{2})^2}{2} \sin^{-1} \frac{4\sqrt{2}}{4\sqrt{2}} \\ &\quad - \frac{4}{2} \sqrt{(4\sqrt{2})^2 - (4)^2} - \frac{(4\sqrt{2})^2}{2} \sin^{-1} \frac{4}{4\sqrt{2}} \\ &= 0 + \frac{16 \times 2}{2} \sin^{-1}(1) - 2\sqrt{32} - 16 - \frac{16 \times 2}{2} \sin^{-1} \left(\frac{1}{\sqrt{2}} \right) \end{aligned}$$


	$= 16 \sin^{-1}(1) - 2\sqrt{16} - 16 \sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$ $= 16 \left[\sin^{-1}(1) - \sin^{-1}\left(\frac{1}{\sqrt{2}}\right) \right] - 8$ $= 16 \left[\frac{\pi}{2} - \frac{\pi}{4} \right] - 8$ $= 16 \left[\frac{4\pi - 2\pi}{4 \times 2} \right] - 8$ $= \frac{16}{8} [2\pi] - 8$ $= 2[2\pi] - 8$ $= 4\pi - 8$ <p>Putting the value of I_1 & I_2 in (1)</p> $\text{Area} = 8 + 4\pi - 8$ $= 4\pi$ <p>\therefore Required Area = 4π Square units</p>	
<p>32</p>	$f'(x) = x^3 - 3x^2 - 10x + 24 = (x-2)(x-4)(x+3) = 0$ $\Rightarrow x = -3, 2, 4$ <p>Strictly $\uparrow = (-3, 2) \cup (4, \infty)$</p> <p>Strictly $\downarrow = (-\infty, 3) \cup (2, 4)$</p> 	<p>3</p>
<p>33</p>	<p>Putting $\sin x = t \Rightarrow \int \frac{2dt}{(1-t)(1+t^2)}$</p> $\frac{2}{(1-t)(1+t^2)} = \frac{A}{1-t} + \frac{Bt+C}{1+t^2}$ <p>$A=1, B=1, C=1$</p> $\int \frac{1}{1-t} dt + \frac{1}{2} \int \frac{2t}{1+t^2} dt + \int \frac{1}{1+t^2} dt \Rightarrow -\log(1-\sin x) + \frac{1}{2} \log(1+\sin^2 x) + \tan^{-1}(\sin x) + C$ <p>OR</p> $\frac{1}{(x+2)(x^2+1)} = \frac{A}{x+2} + \frac{Bx+C}{x^2+1}$	<p>3</p> <p>$\frac{1}{2}$</p>

	Getting A=1/5,B=-1/5,C=2/5 Finally a=-1/10 ,b= -2/10	1+1/2 1
34	$Xdy-ydx=\sqrt{x^2+y^2} dx$ $\Rightarrow Xdy=(\sqrt{x^2+y^2}+y)dx$ $\Rightarrow \frac{dy}{dx} = \frac{(\sqrt{x^2+y^2}+y)}{x}$ homogeneous differential equation $y=v(x)$ $\Rightarrow v+x\frac{dv}{dx}=\frac{(\sqrt{x^2+v^2x^2+vx}}{x} =\sqrt{1+v^2} + v$ $\Rightarrow x\frac{dv}{dx} = \sqrt{1+v^2}$ $\Rightarrow \int \frac{dv}{\sqrt{1+v^2}} = \int \frac{dx}{x}$ $\Rightarrow \text{Log} \sqrt{1+v^2} + v = \text{log}x + \text{log}C$ $\Rightarrow \sqrt{1+v^2} + v = Cx$ $\Rightarrow \frac{y}{x} + \sqrt{1 + \frac{y^2}{x^2}} = Cx$	
35	$\log(x^2+y^2) = 2\tan^{-1}\left(\frac{y}{x}\right)$ $\frac{1}{x^2+y^2} \left[2x + 2y \frac{dy}{dx} \right] = \frac{2}{1+\frac{y^2}{x^2}} \left[\frac{x\frac{dy}{dx}-y.1}{x^2} \right]$ $\Rightarrow \frac{x+y\frac{dy}{dx}}{x^2+y^2} = \frac{x^2}{x^2+y^2} \left[\frac{x\frac{dy}{dx}-y.1}{x^2} \right]$ $\Rightarrow x+y\frac{dy}{dx} = x\frac{dy}{dx} - y \Rightarrow \frac{dy}{dx} = \frac{x+y}{x-y}$	
SECTION V		
36	$\begin{bmatrix} 6 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{bmatrix} = AB = 6I \Rightarrow A^{-1} = \frac{1}{6}B$ $\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 12 \\ -6 \\ 24 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}$ <p>OR</p> $3x+0y+3z=8+2y \Rightarrow 3x-2y+3z=8$ $2x+1y+0z=1+z \Rightarrow 2x+y-z=1$ $4x+0y+2z=4+3y \Rightarrow 4x-3y+2z=4$ <p>Ax=B</p> $\Rightarrow \begin{bmatrix} 3 & 2 & -3 \\ 2 & 1 & -1 \\ 4 & -3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 8 \\ 1 \\ 4 \end{bmatrix}$ $A^{-1} = \frac{1}{17} \begin{bmatrix} 1 & 5 & 1 \\ 8 & 6 & -9 \\ 10 & -1 & 7 \end{bmatrix}$ $X=A^{-1}B \Rightarrow X = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$	<p>5</p> <p>1.5</p> <p>½</p> <p>1.5</p> <p>1.5</p>

37

First we will convert the given inequations into equations, we obtain the following equations:
 $3x+2y=80, 2x+3y=70, x=0$ and $y=0$

5



The corner points of the feasible region are $O(0, 0)$, $A(\frac{80}{3}, 0)$, $E(20, 10)$ and $D(0, \frac{70}{3})$.

The values of Z at these corner points are as follows.

Corner point	$Z = 15x + 10y$
$O(0, 0)$	$15 \times 0 + 10 \times 0 = 0$
$A(\frac{80}{3}, 0)$	$15 \times \frac{80}{3} + 10 \times 0 = 400$
$E(20, 10)$	$15 \times 20 + 10 \times 10 = 400$
$D(0, \frac{70}{3})$	$15 \times 0 + 10 \times \frac{70}{3} = \frac{700}{3}$

We see that the maximum value of the objective function Z is 400 which is at $A(\frac{80}{3}, 0)$ and $E(20, 10)$.
 Thus, the optimal value of Z is 400.

OR

Plane $= 0x + 3y + 4z = 6$

$\vec{x} = 0\hat{i} + 3\hat{j} + 4\hat{k}$

$\vec{OP} = x\hat{i} + y\hat{j} + z\hat{k}$

$\frac{x_1}{0} = \frac{y_1}{3} = \frac{z_1}{4} = r \Rightarrow x_1 = 0, y_1 = 3r, z_1 = 4r$

(x, y, z) lies on plane $\Rightarrow 0 + 3(3r) + 4(4r) = 6$

$\Rightarrow r = 6/25$

Foot of perpendicular $= (0, 18/25, 24/25)$

