

**MARKING SCHEME**

**SAMPLE PAPER 1**

**SECTION A**


Q.No.	Value Point	Marks
1(i)	D	1
(ii)	B OR A	1
(iii)	A	1
(iv)	C	1
2(i)	B	1
(ii)	A	1
(iii)	A	1
(iv)	A OR B	1
3	C	1
4	D OR C	1
5	C	1
6	B OR B	1
7	B OR D	1
8	A OR A	1
9	C	1
10	A	1
11	A	1
12	A	1
13	D	1
14	B OR B	1
15	B	1
16	A	1







**QB365 - Question Bank Software**




	<b>OR</b>	
	<p>Ans : let the rate law expression be <math>\text{Rate} = k [\text{P}]^x [\text{Q}]^y</math>                      from the table we know that  <math>\text{Rate 1} = 3.0 \times 10^{-4} = k (0.10)^x (0.10)^y</math>  <math>\text{Rate 2} = 9.0 \times 10^{-4} = k (0.30)^x (0.30)^y</math>  <math>\text{Rate 3} = 3.0 \times 10^{-4} = k (0.10)^x (0.30)^y</math></p> <p><math>\text{Rate 1} / \text{Rate 3} = (1/3)^y</math> or <math>1 = (1/3)^y</math>                      So <math>y = 0</math>  <math>\text{Rate 2} / \text{Rate 3} = (3)^x</math> or <math>3 = (3)^x</math>                      So <math>x = 1</math>  <math>\text{Rate} = k [\text{P}]</math></p>	<p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p>1</p>
21	<p><math>k = 0.693/t_{1/2}</math>  <math>k = 0.693/5730 \text{ years}^{-1}</math>  <math>t = \frac{2.303 \log C_0}{k C_t}</math>                      let <math>C_0 = 1</math> <math>C_t = 3/10</math> so <math>C_0/C_t = 1 / (3/10) = 10/3</math>  <math>t = \frac{2.303 \times 5730 \log \frac{10}{3}}{0.693}</math>  <math>t = 19042 \times (1 - 0.4771) = 9957 \text{ years}</math></p>	<p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>
22	<p> <math>\text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \underset{\text{OH}}{\text{CH}} - \text{CH}_3 \xrightarrow{\text{H}^+} \text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \underset{\text{OH}_2^+}{\text{CH}} - \text{CH}_3</math>  <math>\text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \underset{\text{OH}_2^+}{\text{CH}} - \text{CH}_3 \xrightarrow{-\text{H}_2\text{O}} \text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \overset{+}{\text{C}}\text{H} - \text{CH}_3</math>  <math>\text{CH}_3 - \overset{+}{\text{C}}\text{H} - \text{CH}_3 \xrightarrow{1,2\text{-hydride shift}} \text{CH}_3 - \overset{+}{\text{C}}(\text{CH}_3) - \text{CH}_2 - \text{CH}_3</math>  <math>\text{CH}_3 - \overset{+}{\text{C}}(\text{CH}_3) - \text{CH}_2 - \text{CH}_3 + \text{Br}^- \rightarrow \text{CH}_3 - \overset{\text{Br}}{\text{C}}(\text{CH}_3) - \text{CH}_2 - \text{CH}_3</math> </p>	<p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>
23	<p><math>\text{XeF}_6</math></p> <p>Central atom Xe has 8 valence electrons, it forms 6 bonds with F and has 1 lone pair. According to VSEPR theory, presence of 6 bp and 1 lp results in distorted octahedral geometry</p>	<p>1</p> <p>1</p>





**QB365 - Question Bank Software**


		
24.	<p>Racemic mixture will be given by 2 chlorobutane as it is an optically active compound.</p> <p>When 2 chlorobutane undergoes <math>S_N^1</math> reaction, both front and rear attack are possible, resulting in a racemic mixture</p>	1 1
25	<p>Let no. of Atoms of element P be <math>x</math></p> <p>No. of tetrahedral voids = <math>2x</math></p> <p>No. Of octahedral voids = <math>x</math></p> <p>Atoms of Q = <math>\frac{1}{3}(2x) + x = \frac{5x}{3}</math></p> <p><math>P_xQ_{5x/3}</math></p> <p><math>P_3Q_5</math></p>	1/2  1/2  1
<b>SECTION C</b>		
26	<p>(i) Due to large surface area and ability to show variable oxidation states</p> <p>(ii) Due to high value of third ionisation enthalpy</p> <p>(iii) Mo(VI) and W(VI) are more stable than Cr(VI).</p> <p align="center"><b>OR</b></p> <p>(i) The general trend towards less negative <math>E^\circ</math> V values across the series is related to the general increase in the sum of the first and second ionisation enthalpies.</p> <p>(ii) The high energy to transform Cu(s) to <math>Cu^{2+}</math> (aq) is not balanced by its hydration enthalpy.</p> <p>(iii) The stability of the half-filled <math>d</math> sub-shell in <math>Mn^{2+}</math> and the completely filled <math>d^{10}</math> configuration in <math>Zn^{2+}</math> are related to their more negative <math>E^\circ</math> V values</p>	1 1 1  1 1 1  1
27	<p>(i) Aniline, <i>N</i>-ethylethanamine, Etanamine</p> <p>(ii) Ethanamine, ethanol, ethanoic acid</p> <p>(iii) <i>N,N</i> dimethylmethanamine, methanamine, <i>N</i>-methylmethanamine</p> <p align="center"><b>OR</b></p> <p>(i) <i>N</i>-methyl ethanamine is a secondary amine. When it reacts with benzenesulphonyl chloride, it forms <i>N</i>- Ethyl -<i>N</i> methyl sulphonamide while and</p>	1 1 1  1

**QB365 - Question Bank Software**

	<p>N,N-dimethyl ethanamine is a tertiary amine it does not react with benzenesulphonyl chloride.</p> <p>(ii) </p> <p>(iii) Butan-1-ol Alcohol forms stronger hydrogen bonds with water than formed by amine due to higher electronegativity of O in alcohol than N in amine</p>	<p>1</p> <p>1/2</p> <p>1/2</p>
28	<p>We know that <math>d = zM / N_a a^3</math></p> <p>For fcc, <math>z=4</math> therefore <math>d = 4 \times M / N_a (3.5 \times 10^{-8})^3 \text{ g/cm}^3</math></p> <p>For bcc, <math>z=2</math> therefore <math>d = 2 \times M / N_a (3.0 \times 10^{-8})^3 \text{ g/cm}^3</math></p> <p><math>d/d' = 4/(3.5 \times 10^{-8})^3 / 2/(3.0 \times 10^{-8})^3 = 1.26:1</math></p>	<p>1/2</p> <p>1</p> <p>1</p> <p>1/2</p>
29	<p>(i) </p> <p></p> <p>(ii) </p>	<p>1</p> <p>1</p> <p>1</p>
30	<p>i. Arrange the following in decreasing order of bond dissociation enthalpy <math>\text{Cl}_2 &gt; \text{Br}_2 &gt; \text{F}_2 &gt; \text{I}_2</math></p> <p>ii. Bi does not form <math>p\pi-p\pi</math> bonds as its atomic orbitals are large and diffuse so effective overlapping is not possible</p> <p>iii. Due to small size of oxygen, it has greater electron electron repulsions</p>	<p>1</p> <p>1</p> <p>1</p>
<b>SECTION D</b>		
31.	<p>(i) (a) <math>3\text{Cu} + 8 \text{HNO}_3(\text{dilute}) \rightarrow 3\text{Cu}(\text{NO}_3)_2 + 2\text{NO} + 4\text{H}_2\text{O}</math> (b)</p>	<p>1</p>

	<div style="text-align: center;">  </div> <p>(ii) 'X' is Helium                      It is used as a diluent for oxygen in modern diving apparatus because of its very low solubility in blood.                      It monoatomic having no interatomic forces except weak dispersion forces and has second lowest mass therefore bp is lowest.</p> <p style="text-align: center;">OR</p> <p>(a) <math>\text{H}_2\text{Te}</math>, <math>\text{H}_2\text{Se}</math>, <math>\text{H}_2\text{S}</math>, <math>\text{H}_2\text{O}</math>                      (b) <math>[\text{Fe}(\text{H}_2\text{O})_5(\text{NO})]^{2+}</math></p> <p>(ii) A is chlorine gas                      Its bleaching action is due to oxidation.  <math>\text{Cl}_2 + \text{H}_2\text{O} \rightarrow 2\text{HCl} + \text{O}</math>, Coloured substance + O <math>\rightarrow</math> Colourless substance  <math>6\text{NaOH} + 3\text{Cl}_2 \rightarrow 5\text{NaCl} + \text{NaClO}_3 + 3\text{H}_2\text{O}</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
<p>32</p>	<div style="text-align: center;"> <p>A: <math>\text{C}\equiv\text{CH}</math>      B: <math>\text{COCH}_3</math>      C: <math>\text{COOK}</math>      D: <math>\text{CHI}_3</math></p>     </div>	<p><math>\frac{1}{2}</math> each</p> <p>1</p> <p>1</p>

	<p>  </p> <p>  </p> <p> <b>OR</b> </p> <p>           (i) <math>\text{CH}_3\text{COCH}_3 + \text{CH}_3\text{CHO}</math>  <math>\downarrow</math> dil NaOH  <math>(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{CHO} + \text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{COCH}_3</math>  <math>\downarrow</math> Heat  <math>(\text{CH}_3)_2\text{C}=\text{CHCHO} + \text{CH}_3\text{CH}=\text{CHCOCH}_3</math> </p> <p>           (ii) (a)  </p> <p>           (b) <math>\text{CH}_3\text{COCH}_3 \xrightarrow{\text{H}_2, \text{Pd}} \text{CH}_3\text{CH}(\text{OH})\text{CH}_3 \xrightarrow{\text{H}_2\text{SO}_4} \text{CH}_3\text{CH}=\text{CH}_2</math> </p> <p>           (c)  </p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
33	<p>(i) limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of the anion and cation of the electrolyte.</p> <p>(ii) <math>E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}} = 0.34 - (-1.66) = 2.00 \text{ V}</math>  <math>E_{\text{cell}} = E^\circ_{\text{cell}} - 0.059 \log [\text{Al}^{3+}]^2</math></p>	<p>1</p> <p>1/2</p>

<p style="text-align: center;"><math>n \quad [\text{Cu}^{2+}]^3</math></p>	1
<p>Here <math>n = 6</math></p>	1/2
$E_{\text{cell}} = 2 - \frac{0.059}{6} \log \frac{[0.15]^2}{[0.025]^3}$	1
$= 2 - 0.059/6 (2 \log 0.15 - 3 \log 0.025)$	1
$= 2 - 0.059/6 (-1.6478 + 4.8062) = 2 - 0.0311 = 1.9689\text{V}$	1
<b>OR</b>	
(i) $\text{MnO}_4^-$	1
(ii)(a) Molar conductivity of a solution at a given concentration is the conductance of the volume $V$ of solution containing one mole of electrolyte kept between two electrodes with area of cross section $A$ and distance of unit length.	1
(b) Strong electrolyte, For strong electrolytes, $\Lambda_m$ increases slowly with dilution	1
(c) $\Lambda_m = \Lambda_m^\circ - A c^{1/2}$	1
Therefore $\Lambda_m^\circ = 150 \text{ S cm}^2 \text{ mol}^{-1}$	1
(d)	1
	1
$A = -\text{slope} = - (149 - 147.8 / 0.010 - 0.022) = 100 \text{ S cm}^2 \text{ mol}^{-1} / (\text{mol/L}^{-1})^{1/2}.$	1